

What can a podiatrist do?

A podiatrist can explain how to reduce the risk of foot blisters, advise on their treatment and check that your feet are generally in good condition. Podiatrists can also provide expert advice on the use of insoles (also known as orthotics) and other appliances to keep your feet healthy and comfortable.

How to Contact a Podiatrist

To find a local Health Professionals Council (HPC) registered podiatrist, you can either approach your GP practice for information on an NHS referral or if you decide to seek a private podiatrist you could use the following link on the Society's

website <http://www.feetforlife.org/cgi-site/chiropracist.cgi> or look at the front of the chiropracists and podiatrists section in Yellow Pages for The Society of Chiropracists and Podiatrists listing.

Further Foot Health Information

The Society of Chiropracists and Podiatrists provides a number of foot health information leaflets as either PDFs to download or in hard copy form, please see the Society's website, www.feetforlife.org for further information.

Useful Contacts

www.feetforlife.org
www.hpc-uk.org

With the compliments of:

The Society of Chiropracists and Podiatrists,
1 Fellmongers Path, Tower Bridge Road, London, SE1 3LY,
Tel: 0845 450 3720 • Fax: 0845 450 3721
Email: enq@scpod.org • www.feetforlife.org

Copyright © 2008 The Society of Chiropracists and Podiatrists
Designed and printed by: Harvest Digital Imaging Ltd



The Society of Chiropracists and Podiatrists

Blisters



The Society of
Chiropracists and
Podiatrists

www.feetforlife.org

Blisters



Blisters are painful, fluid-filled lesions, often caused by friction and pressure. The most common cause is repeated rubbing from ill-fitting footwear and socks, and excessive moisture between the foot and the sock.

Blisters can also be caused by foot deformities, and be of concern to those with existing conditions such as Diabetes. If you suffer from one of these conditions please refer to your podiatrist for advice. To find a podiatrist in your area please turn to the back page of this leaflet.

How to Prevent Blisters

As blisters are a result of friction there are a number of simple techniques that can prevent your walking, running or social activity being ruined by a blister:

- Wear comfortable, good-fitting footwear especially on long walks or runs.

- Wear good walking socks in the right size, look for ones made of fibres with good 'wicking' properties. Try wearing socks inside out to prevent the seams rubbing. Change your socks daily!
- Some walkers and runners wear special 'dual layer' socks. The inner layer moves with the foot, the outer with the shoe - eliminating friction at the skin surface. Ensure you change to a fresh pair after they become damp.
- Keep your toenails trim
- If your feet get very sweaty there are products that can help control excessive moisture. Your podiatrist can tell you about these.
- Quickly remove any foreign bodies from your socks and boots

- Ensure that the tongue and laces of your boots are arranged correctly and evenly
- Check your feet carefully and regularly for any sign of rubbing and tenderness

If you feel a blister developing do this:

Act immediately you feel any friction or discomfort: blisters can form very quickly. If you feel a blister developing, stop walking, take your boots and socks off and examine your feet. Consider applying some material cushioning or padding, or a breathable waterproof plaster.



If a blister has developed

Most foot blisters filled with clear fluid last between three and seven days, and will normally clear up if further excessive friction is avoided. If a blister does occur, do not pop it. Cut a hole in a piece of foam or felt, forming a 'doughnut' over the blister; tape the foam or felt in place or cover with a soft gel-type dressing. Treat an open blister with mild soap and water; cover it with an anti-septic ointment and protective soft gel dressing to prevent infection and speed up the healing process.

